



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MORTALITY—FOREIGN AND INSULAR—COUNTRIES AND CITIES
(untabulated).

ARGENTINA—*Buenos Aires*.—Month of August, 1910. Population 1,275,910. Total number of deaths from all causes 2,048, including smallpox 29, measles 6, scarlet fever 3, diphtheria 16, tuberculosis, 247.

AZORES—*St. Michaels*.—Month of August, 1910. Population 131,183. Total number of deaths from all causes 110, including typhoid fever 1, tuberculosis 6.

CANADA—*Sherbrooke*.—Month of October, 1910. Population 15,300. Total number of deaths from all causes 20, including diphtheria 1, tuberculosis 4.

CHILE—*Punta Arenas*.—Month of August, 1910. Population 12,000. Total number of deaths from all causes 43, including tuberculosis 3.

EAST AFRICA—*Lourenco Marquez*.—Month of September, 1910. Population, 10,000. Total number of deaths from all causes 58, including tuberculosis 17.

FRANCE—*Nice*.—Month of July, 1910. Population 150,881. Total number of deaths from all causes 190, including typhoid fever 1, measles 1, tuberculosis 29.

Month of August, 1910. Total number of deaths 157, including typhoid fever 4, diphtheria 1, tuberculosis 22.

GREAT BRITAIN.—Week ended October 15, 1910.

England and Wales.—The deaths registered in 77 great towns correspond to an annual rate of 13.1 per 1,000 of the population, which is estimated at 16,940,895.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 16.9 per 1,000 of the population, which is estimated at 1,151,790. The lowest rate was recorded at Drogheda, viz, 4.1, and the highest at Clonmel, viz, 41 per 1,000.

Scotland.—The deaths registered in 8 principal towns correspond to an annual rate of 12.9 per 1,000 of the population, which is estimated at 1,865,571. The lowest rate was recorded at Aberdeen, viz, 9.8, and the highest at Dundee, viz, 17.5. The total number of deaths from all causes was 468, including typhoid fever 2, measles 3, scarlet fever 4, diphtheria 14.

ITALY—*Genoa*.—Two weeks ended October 31, 1910. Population 279,163. Total number of deaths from all causes 187, including typhoid fever 3, tuberculosis 20.

SPAIN—*Almeria*.—Month of October, 1910. Population 50,910. Total number of deaths from all causes 116, including typhoid fever 2, scarlet fever 1, tuberculosis 11.

Seville.—Month ended October 31, 1910. Population 154,315. Total number of deaths from all causes 384, including smallpox 2, scarlet fever 4, diphtheria 9, tuberculosis 67.

By authority of the Secretary of the Treasury:

WALTER WYMAN,

Surgeon-General,

United States Public Health and Marine-Hospital Service.